


LOW ENERGY KAON PHYSICS AT DAΦNE

P. GIANOTTI

*INFN National Laboratory of Frascati, Via E. Fermi 40,
I-00044 Frascati, Italy*




DAΦNE e^+e^- collider is an abundant source of low energy $K\bar{K}$ pairs suitable to explore different fields of non perturbative QCD regime. Two different experiments, DEAR and FINUDA, using different experimental techniques are trying to shed new light on the strong interaction at the nucleon scale by producing high precision results at this energy range. The DEAR experiment is studying kaonic atoms in order to determine antikaon-nucleon scattering lengths. FINUDA aims to produce hypernuclei to study nuclear structure and Λ -N interaction.

1 The DAΦNE Φ -factory

DAΦNE¹ is a positron/electron collider working at the energy of the Φ resonance (1.02 GeV c.m.). This resonance decays mainly into low energy (~ 16 MeV) $K\bar{K}$ pairs that can be used to perform different physics studies. The DAΦNE machine has two interactions regions (IR1, IR2) where three experiments take place: the KLOE one is installed in IR1, DEAR and FINUDA experiments are sharing IR2.

DAΦNE has been designed to provide high luminosity in order to allow precise studies of rare phenomena such as CP violation in the $K\bar{K}$ system, and rare kaon decay studies. Furthermore, this characteristic turns out to be very useful to explore many experimental physics fields.

After the commissioning phase, DAΦNE started delivering luminosity to the KLOE and DEAR experiments. Fig. 1 shows the plots of the delivered luminosity in the period 2000-2002 together with some relevant machine design parameters. The DAΦNE performances improved considerably during the last year. At present the collider has reached a peak luminosity of $0.8 \cdot 10^{32}$ while the daily integrated luminosity is $\sim 4 \text{ pb}^{-1}$ allowing the experiments to produce good quality data.



Single beam energy	0.51 GeV
Number of particles per bunch	8.9×10^{10}
Number of bunches per ring	up to 120
Crossing frequency	up to 368.25 MHz
Horizontal emittance	1.0 mm.mrad
Vertical emittance	0.01 mm.mrad
Horizontal beta function at crossing	4.5 m
Vertical beta function at crossing	0.045 m
Total crossing angle in the horizontal plane	20 to 30 mrad
Horizontal beam-beam tune shift per	0.04
Vertical beam-beam tune shift per crossing	0.04
Bunch length	30 mm r.m.s.
Horizontal beam size at crossing	2.0 mm r.m.s.
Vertical beam size at crossing	0.02 mm r.m.s.
Maximum stored current per ring	5.2 A
Maximum luminosity	$5.3 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$

Figure 1: Plots of the luminosity delivered by DAΦNE to the experiments in the period 2000-2002. The table on the right shows some relevant machine parameters.

2 The DEAR experiment

DEAR (DAΦNE Exotic Atom Research) ² aims to evaluate precisely the isospin dependent antikaon-nucleon scattering lengths. This is done by measuring the strong interaction shifts and widths of the x-ray lines emitted in the transitions to the ground levels of kaonic hydrogen and kaonic deuterium atoms ⁴. When a negative kaon enters the target, it can lose some energy through ionization and excitation processes of the atoms and molecules of the medium. When its kinetic energy is very low, it can eventually form a hydrogen-like atom by replacing an electron in an excited state. The newly formed atom can deexcite by means of three different processes: dissociation of the surrounding molecules, external Auger transitions, and radiative transitions. When the kaon reaches the low-n states, it suffers strong interactions with the nucleus which produce a shift of the energy of the low-lying levels from the pure electromagnetic values. On the other hand, the decreased lifetime of the levels result in an increase of the observed levels width. The shift ϵ and the width Γ of the 1s state of kaonic hydrogen/deuterium are related in a fairly model-independent way to the isospin dependent scattering lengths. The measurement performed by DEAR is then crucial, since an accurate determination of these parameters will define better the K^- -N dynamics placing strong constraints to the SU(3) description of the chiral symmetry breaking (see ³ for more theoretical details).

2.1 The experimental setup

The scientific program of DEAR consists of two phases. During phase one, the cryogenic target, is filled with pure nitrogen at 85 K resulting in a density of about 4.4 times the NTP one. The purposes of these preliminary kaonic nitrogen measurements are the background level evaluation and reduction, the test of the reliability of the MonteCarlo simulation code, and the tuning of the degrader thickness placed at the entrance of the target. Nitrogen has been chosen in order to have a faster feedback on the detector capabilities since the yield of the signal of kaonic nitrogen is almost 10 times greater than that of kaonic hydrogen.

When the detector setup has been optimized the experiment moves to phase two to perform the measurements on kaonic hydrogen. The target temperature has been set to 23 K and the pressure to 1.82 bar. At these working conditions the hydrogen density is 3.1% of liquid

hydrogen. This setup has been chosen as a compromise to enhance the atom production yield without destroying the x-ray signal.

To detect the x rays with high resolution and efficiency, 16 CCD detectors are placed all around the target. The CCDs are essentially silicon-based MOS devices with a pixel structure. CCDs are ideal for a wide range of applications in the energy range from few eV up to 20 KeV because, apart from high energy resolution, they provide a very good charged particles and γ 's background rejection ($\sim 10^{-5}$). DEAR CCDs have a thermal noise FWHM of 15 eV, and an energy resolution at 5.9 KeV (Mn K_{α} line) of 136 eV FWHM. CCDs cannot be triggered and an x-ray signal is recognized in a topological way as an isolated hit pixel. On the contrary, the energy lost by charge particles, neutrons and gammas is normally distributed over several pixels.

2.2 Preliminary results

During year 2002 DEAR had two run periods: the first one was performed with the nitrogen target setup in order to study and reduce the background, and to measure kaonic nitrogen; the second was dedicated to the measurements on kaonic hydrogen.

At DAΦNE the main source of background is the Touschek effect, an elastic scattering among the particles of the same bunch. This effect produce a broadening of the momentum distribution of the bunch making some particles of the beam to go out of the machine acceptance. The CCD detectors cannot be triggered, and thus the lost particles can disturb the measurements. Great efforts have been done to tune the machine parameters to reduce this background, and furthermore, different shielding materials have been placed all around the DEAR detector. Nevertheless, a high background is still present. Fig. 2 shows the x-ray spectrum obtained integrating 10 pb^{-1} on the nitrogen target without (*left*) and with (*right*) the background subtraction. Several x-ray lines are clearly observed: the two lines at 4.6 KeV and 7.6 KeV correspond to kaonic nitrogen transitions; the other ones are due to electronic x-ray transitions inside the materials present in the setup. This extra lines are useful to calibrate the absolute energy scale.




Figure 2: X-ray spectra obtained with the NTP nitrogen setup. (*left*) Complete spectrum; (*right*) background subtracted spectrum.

Fig. 3 is the preliminary result of the analysis of 20 pb^{-1} taken with the hydrogen setup. In the background subtracted spectrum (*right*) are visible hints of kaonic hydrogen x-rays transitions.

At present, the data analysis of all the 58 pb^{-1} taken with hydrogen is in progress. Pure background data are also under analysis in order to eliminate from the spectrum any spurious signal not coming from kaonic hydrogen. The expected precision to be obtained on the shift of the K_{α} line of kaonic hydrogen is around 15%.




Figure 3: X-ray spectra obtained with the cryogenic hydrogen setup. (left) Complete spectrum; (right) background subtracted spectrum.

The experiment long term plans foresee to replace the CCDs with silicon drift detectors. These devices have a similar energy resolution, and moreover can be triggered, allowing an on-line background reduction. With the new detector setup a new set of measurements with nitrogen, hydrogen and deuterium will be performed in the next years. The goal is to reach a percent level precision on antikaon-proton and antikaon-neutron scattering lengths.

3 The FINUDA experiment

On January 2003 the DEAR detector has been moved out from DAΦNE IR2 and a new experiment FINUDA (Fisica Nucleare a DAΦNE) has been installed in place of it. FINUDA⁵ will start the data taking during the summer of 2003 in order to study ΛN interactions by means of hypernuclei. An hypernucleus is a nuclear system in which a neutron has been replaced by a Λ hyperon. At DAΦNE this will be obtained by stopping the negative kaons from the Φ decay into a thin (~ 0.1 g/cm²) nuclear target. In the past hypernuclear physics has probably reached its best result in proving the validity of the nuclear shell model not only on the nuclear surface, but also deeply inside the nuclear matter. This allowed to elaborate combined precise models of $N - N$ and $\Lambda - N$ interactions for which today the main terms are known with good precision. On the other hand, spin-spin and spin-orbit contributions are still not so well determined, and new high quality data can help the work of the theoreticians. Furthermore, the investigation of non-mesonic hypernuclear decays, $\Lambda N \rightarrow NN$, allow to study the four-fermions, strangeness-changing weak process, hardly achievable with other experimental techniques. Whether the $\Delta I = 1/2$ rule, which governs free Λ decay, also applies to the non-mesonic decays remain an open problem. For a recent review of the main topics related to the hypernuclear physics field see ref.⁶

1. M. Zobov *et al*, Proceeding of EPAC2002, Paris, France.
2. S. Bianco *et al*, *Rivista del Nuovo Cimento* **22** No. 11, 1 (1999).
3. G. Colangelo, this proceeding.
4. C. Guaraldo, Proceeding of *PHYSICS AND DETECTORS FOR DAΦNE*, Frascati Physics Series **Vol. XVI**, 643 (1999).
5. P. Gianotti, *Nucl. Phys.* **A691**, 483c (2001).
6. Proceeding of the *VII International Conference on Hypernuclear and Strange Particle Physics*, *Nucl. Phys.* **A691** (2001).

This figure "foto_paola.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/nucl-ex/0305010v1>